## What is claimed is:

1

2

1

2

3

the first that

: Pa

3

1

2

3

1

2

1

2

- 1. A method of synthesizing carbon nanotubes, comprising the steps of: introducing a catalyst in a reactor; supplying a reactant gas containing a carbon source gas over the catalyst; selectively and locally heating the catalyst in the reactor; and growing carbon nanotubes from the heated catalyst.
- 2. The method of claim 1, wherein the catalyst is formed of a transition metal such as iron, nickel or cobalt; metal sulfide, metal carbide, metal oxide or metal salt of the transition metal; or an organic compound containing the transition metal.
- 3. The method of claim 1, wherein the catalyst is loaded on a support by an impregnation method, an incipient wetness method or an ion-exchange method and is supplied into the reactor in a powder state.
- 4. The method of claim 1, wherein the catalyst is loaded on a substrate by a deposition method, a painting method and a spray method to be supplied into the reactor.
- 5. The method of claim 1, wherein for the catalyst, a metal precursor is loaded on a substrate or a support and changed into a metal phase through reduction, calcination, sulfiding or carbonization, and the metal catalyst is supplied into the reactor.
- 6. The method of claim 1, wherein for the catalyst, metal sulfide obtained by sulfiding a metal precursor with hydrogen sulfide is used.
- 7. The method of claim 1, wherein the catalyst is supplied into the reactor in the form of a catalyst precursor in gas phase.

1

2

1

2

5

1

1

2

- 8. The method of claim 7, wherein the catalyst precursor is ferrocene or 1 iron pentacarbonyl. 2
- 9. The method of claim 1, wherein the carbon source gas contains one selected from the group consisting of acetylene, methane, propane and benzene. 2
  - 10. The method of claim 1, wherein the reactant gas further comprises hydrogen gas or inert gas.
  - 11. The method of claim 1, wherein the reactant gas further comprises hydrogen sulfide (H<sub>2</sub>S) gas.
  - 12. The method of claim 1, wherein the local heating of the catalyst is performed by irradiation of microwaves.
  - 13. The method of claim 1, wherein the local heating of the catalyst is performed by electromagnetic inductive heating.
  - 14. The method of claim 1, wherein the local heating of the catalyst is performed by laser heating.
  - 15. The method of claim 1, wherein the local heating of the catalyst is performed by radio frequency heating.
  - 16. An apparatus for synthesizing carbon nanotubes, comprising: a reactor for receiving a catalyst;
- a reactant gas supplier for supplying a carbon source gas into the reactor; 3 and 4
  - a local heater for selectively heating the catalyst received in the reactor.